

SAMPLE CONTENT

33
YEARS

1988
to
2020



NEET CHEMISTRY

PSP PREVIOUS
SOLVED
PAPERS

● TOPIC-WISE AND SUBTOPIC-WISE ●

Includes Solved Questions of 2020

A comprehensive collection of NEET & AIPMT Questions from past 33 Years

1571 MCQs

★ In accordance with 11th and 12th NCERT Books ★

Target Publications[®] Pvt. Ltd.

33
YEARS

1988
to
2020

NEET CHEMISTRY

PSP PREVIOUS
SOLVED
PAPERS

- TOPIC - WISE AND SUBTOPIC - WISE

Salient Features

- ☞ A compilation of 33 years of AIPMT/NEET questions (1988-2020)
- ☞ Includes solved questions from NEET 2020
- ☞ Includes '1571' AIPMT/NEET MCQs
- ☞ Topic - wise and Subtopic - wise segregation of questions
- ☞ Year-wise flow of content concludes with the latest questions
- ☞ Relevant solutions provided
- ☞ Graphical analysis of questions – Topic - wise and Subtopic - wise

Printed at: **Print Vision**, Navi Mumbai

© Target Publications Pvt. Ltd.

No part of this book may be reproduced or transmitted in any form or by any means, C.D. ROM/Audio Video Cassettes or electronic, mechanical including photocopying; recording or by any information storage and retrieval system without permission in writing from the Publisher.

TEID: 1673

P.O. No. 1195

PREFACE

Target's 'NEET: Chemistry PSP (Previous Solved Papers)' is a compilation of questions asked in the past 33 years (1988-2020) in the National Eligibility cum Entrance Test (NEET), formerly known as the All India Pre-Medical Test (AIPMT). The book is crafted in accordance with the Std. XI and Std. XII NCERT textbook.

The book consists of topic - wise categorization of questions. Each chapter is further segregated into subtopics and thereafter all the questions pertaining to a subtopic are arranged year-wise starting with the latest year. To aid students, we have also provided hints for questions wherever deemed necessary.

A graphical (% wise) analysis of the subtopics for the past 33 years as well as 8 years (2013 onwards) has been provided at the onset of every topic. Both the graphs will help the students to understand and analyse each subtopic's distribution for AIPMT (33 years) and NEET-UG (8 Years).

We are confident that this book will comprehensively cater to needs of students and effectively assist them to achieve their goal.

We welcome readers' comments and suggestions which will enable us to refine and enrich this book further.

All the best to all Aspirants!

Yours faithfully,
Authors
Edition: Second

Frequently Asked Questions

Why this book?	<ul style="list-style-type: none"> This book acts as a go-to tool to find all the AIPMT/NEET questions since the past 33 years at one place. The subtopic wise arrangement of questions provides the break-down of a chapter into its important components which will enable student to design an effective learning plan. The graphical analysis guides students in ascertaining their own preparation of a particular topic.
Why the need for two graphs?	<p>Admission for undergraduate and post graduate medical courses underwent a critical change with the introduction of NEET in 2013. Although it received a huge backlash and was criticised for the following two years, NEET went on to replace AIPMT in 2016. The introduction of NEET brought in a few structural differences in terms of how the exam was conducted. Although the syllabus has largely remained the same, the chances of asking a question from a particular subtopic is seen to vary slightly with the inception of NEET.</p> <p>The two graphs will fundamentally help the students to understand that the (weightage) distribution of a particular topic can vary i.e., a particular subtopic having the most weightage for AIPMT may not necessarily be the subtopic with the most weightage for NEET.</p>
How are the two graphs beneficial to the student?	<ul style="list-style-type: none"> The two graphs provide a subtopic's weightage distribution over the past 33 years (for AIPMT) and over the past 8 years (for NEET-UG). The students can use these graphs as a self-evaluation tool by analyzing and comparing a particular subtopic's weightage with their preparation of the subtopic. This exercise would help the students to get a clear picture about their strength and weakness based on the subtopics. Students can also use the graphs as a source to know the most important as well as least important subtopics as per weightage of a particular topic which will further help them in planning the study structure of a particular chapter. <i>(Note: The percentage-wise weightage analysis of subtopics is solely for the knowledge of students and does not guarantee questions from subtopics having the most weightage, in the future exams. Question classification of a subtopic is done as per the authors' discretion and may vary with respect to another individual.)</i>

Disclaimer


Utmost care has been taken in compiling and checking the information to ensure that the content is useful and accurate. However, the publisher and the authors shall not be responsible for any loss or damages caused to any person on account of errors or omission which might have crept in or disagreement of any third party on the point of view expressed in the reference book.

© reserved with the Publisher for all the contents created by our Authors.

Index

No.	Topic Name	Page No.
1	Some Basic Concepts of Chemistry	1
2	Structure of Atom	8
3	Classification of Elements and Periodicity in Properties	16
4	Chemical Bonding and Molecular Structure	21
5	States of Matter: Gases and Liquids	31
6	Thermodynamics	38
7	Equilibrium	51
8	Redox Reactions	67
9	Hydrogen	70
10	s-Block Elements	73
11	p-Block Elements	79
12	Organic Chemistry – Some Basic Principles and Techniques	84
13	Hydrocarbons	93
14	Environmental Chemistry	110
15	The Solid State	112
16	Solutions	119
17	Electrochemistry	129
18	Chemical Kinetics	139
19	Surface Chemistry	149
20	General Principles and Processes of Isolation of Elements	152
21	p-Block Elements (Group 15 to 18)	155
22	d and f-Block Elements	165
23	Coordination Compounds	174
24	Haloalkanes and Haloarenes	188
25	Alcohols, Phenols and Ethers	197
26	Aldehydes, Ketones and Carboxylic Acids	208
27	Organic Compounds Containing Nitrogen	224
28	Biomolecules	235
29	Polymers	242
30	Chemistry in Everyday Life	247

Topic-wise Weightage Analysis of past 8 Years (2013 Onwards)




Total No. of Questions: 480

1


Some Basic Concepts of Chemistry

- 1.1 Units of measurement
- 1.2 Uncertainty in measurement
- 1.3 Atomic and molecular masses
- 1.4 Mole concept and molar mass
- 1.5 Percentage composition, empirical and molecular formulae
- 1.6 Chemical reactions, stoichiometry and calculations based on stoichiometry

**33 Years NEET/AIPMT Analysis
(Percentage-wise weightage of sub-topics)**



**8 Years NEET Analysis (2013 Onwards)
(Percentage-wise weightage of sub-topics)**



[Note: Till date no questions have been asked from subtopics: General introduction – Importance and scope of chemistry, Laws of chemical combination, Dalton's atomic theory: concept of elements, atoms and molecules]

1.1 Units of measurement

1. The dimensions of pressure are the same as that of _____. [1995]
 - (A) force per unit volume
 - (B) energy per unit volume
 - (C) force
 - (D) energy

2. An element, X has the following isotopic composition:
 $^{200}\text{X} : 90\%$ $^{199}\text{X} : 8.0\%$ $^{202}\text{X} : 2.0\%$
 The weighted average atomic mass of the naturally-occurring element X is closest to _____. [2007]
 - (A) 201 amu
 - (B) 202 amu
 - (C) 199 amu
 - (D) 200 amu

1.2 Uncertainty in measurement

1. Given numbers: 161 cm, 0.161 cm, 0.161 cm. The number of significant figures for the three numbers is _____. [1998]
 - (A) 3 and 4 respectively
 - (B) 3, 4 and 4 respectively
 - (C) 3, 4 and 5 respectively
 - (D) 3, 3 and 3 respectively

1.4 Mole concept and molar mass

1.3 Atomic and molecular masses

1. Boron has two stable isotopes, ^{10}B (19%) and ^{11}B (81%). Calculate average at.wt. of boron in the periodic table. [1990]
 - (A) 10.8
 - (B) 10.2
 - (C) 11.2
 - (D) 10.0

1. 1 cc N_2O at NTP contains _____. [1988]
 - (A) $\frac{1.8}{224} \times 10^{22}$ atoms
 - (B) $\frac{6.02}{22400} \times 10^{23}$ molecules
 - (C) $\frac{1.32}{224} \times 10^{23}$ electrons
 - (D) all the above
2. The number of oxygen atoms in 4.4 g of CO_2 is approximately _____. [1990]
 - (A) 1.2×10^{23}
 - (B) 6×10^{22}
 - (C) 6×10^{23}
 - (D) 12×10^{23}



3. The molecular weight of O_2 and SO_2 are 32 and 64 respectively. At $15^\circ C$ and 150 mmHg pressure, one litre of O_2 contains 'N' molecules. The number of molecules in two litres of SO_2 under the same conditions of temperature and pressure will be _____.
[1990]
- (A) N/2 (B) N
(C) 2N (D) 4N
4. The number of moles of oxygen in 1 L of air containing 21 % oxygen by volume, in standard conditions, is _____. [1995]
- (A) 0.186 mol (B) 0.21 mol
(C) 2.10 mol (D) 0.0093 mol
5. 0.24 g of a volatile gas, upon vaporisation, gives 45 mL vapour at NTP. What will be the vapour density of the substance? (Density of $H_2 = 0.089$) [1996]
- (A) 95.93 (B) 59.93
(C) 95.39 (D) 5.993
6. Haemoglobin contains 0.334 % of iron by weight. The molecular weight of haemoglobin is approximately 67200. The number of iron atoms (Atomic weight of Fe is 56) present in one molecule of haemoglobin is _____. [1998]
- (A) 4 (B) 6
(C) 3 (D) 2
7. The number of atoms in 4.25 g of P_4 is approximately _____. [2001]
- (A) 4×10^{23} (B) 6×10^{23}
(C) 1×10^{23} (D) 6×10^{23}
8. Specific volume of cylindrical virus particle is 6.02×10^{-2} cc/g whose radius and length are 7 Å and 10 Å respectively. If $N_A = 6.0 \times 10^{23}$ find molecular weight of virus. [2001]
- (A) 7.4 kg/mol
(B) 1.5×10^4 kg/mol
(C) 0.08 kg/mol
(D) 3.0×10^3 kg/mol
9. Which has maximum molecules? [2002]
- (A) 7 g N_2 (B) 2 g H_2
(C) 16 g NO_2 (D) 16 g O_2
10. The maximum number of molecules is present in _____. [2004]
- (A) 15 L of H_2 gas at STP
(B) 5 L of N_2 gas at STP
(C) 0.5 g of H_2 gas
(D) 10 g of O_2 gas
11. The number of atoms in 0.1 mol of a triatomic gas is ($N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$) _____. [2010]
- (A) 6.026×10^{22} (B) 1.806×10^{23}
(C) 3.600×10^{23} (D) 1.800×10^{22}
12. Which has the maximum number of molecules among the following? [2011]
- (A) 44 g CO_2 (B) 48 g O_3
(C) 8 g H_2 (D) 64 g SO_2
13. Equal masses of H_2 , O_2 and methane have been taken in a container of volume V at temperature $27^\circ C$ in identical conditions. The ratio of the volume of gases $H_2 : O_2 : \text{methane}$ would be _____. [2014]
- (A) 8 : 16 : 1 (B) 16 : 8 : 1
(C) 16 : 1 : 2 (D) 8 : 1 : 2
14. A mixture of gases contains H_2 and O_2 gases in the ratio of 1 : 4 (w/w). What is the molar ratio of the two gases in the mixture? [2015]
- (A) 1 : 1 (B) 2 : 1
(C) 1 : 4 (D) 4 : 1
15. The number of water molecules is maximum in _____. [2015]
- (A) 18 g of water
(B) 18 moles of water
(C) 18 molecules of water
(D) 1.8 g of water
16. If Avogadro number N_A , is changed from $6.022 \times 10^{23} \text{ mol}^{-1}$ to $6.022 \times 10^{20} \text{ mol}^{-1}$, this would change _____. [2015]
- (A) the ratio of chemical species to each other in a balanced equation
(B) the ratio of elements to each other in a compound
(C) the definition of mass in units of grams
(D) the mass of one mole of carbon
17. At S.T.P. the density of CCl_4 vapour in g/L will be nearest to _____. [2016]
- (A) 6.87 (B) 3.42
(C) 10.26 (D) 4.57
18. In which case is the number of molecules of water maximum? [2018]
- (A) 18 mL of water
(B) 0.18 g of water
(C) 0.00224 L of water vapours at 1 atm and 273 K
(D) 10^{-3} mol of water
19. Which one of the followings has maximum number of atoms? [2020]
- (A) 1 g of $Mg_{(s)}$ [Atomic mass of Mg = 24]
(B) 1 g of $O_{2(g)}$ [Atomic mass of O = 16]
(C) 1 g of $Li_{(s)}$ [Atomic mass of Li = 7]
(D) 1 g of $Ag_{(s)}$ [Atomic mass of Ag = 108]



1.5 Percentage composition, empirical and molecular formulae

- Which of the following fertilizers has the highest nitrogen percentage? [1993]
 (A) Ammonium sulphate
 (B) Calcium cyanamide
 (C) Urea
 (D) Ammonium nitrate
- Suppose the elements X and Y combine to form two compounds XY_2 and X_3Y_2 . When 0.1 mole of XY_2 weighs 10 g and 0.05 mole of X_3Y_2 weighs 9 g, the atomic weights of X and Y are _____. [Phase-II 2016]
 (A) 30, 20 (B) 40, 30
 (C) 60, 40 (D) 20, 30
- Percentage of Se in peroxidase anhydrous enzyme is 0.5 % by weight (at. Wt. = 78.4) then minimum molecular weight of peroxidase anhydrous enzyme is _____. [2001]
 (A) 1.568×10^4
 (B) 1.568×10^3
 (C) 15.68
 (D) 2.136×10^4

1.6 Chemical reactions, stoichiometry and calculations based on stoichiometry

- A metal oxide has the formula Z_2O_3 . It can be reduced by hydrogen to give free metal and water. 0.1596 g of the metal oxide requires 6 mg of hydrogen for complete reduction. The atomic weight of the metal is _____. [1982]
 (A) 27.9 (B) 55.8
 (C) 79.8 (D) 119.6
- What is the weight of oxygen required for the complete combustion of 2.8 kg of ethylene? [1989]
 (A) 2.8 kg (B) 6.4 kg
 (C) 9.6 kg (D) 96 kg
- The number of gram molecules of oxygen in 6.02×10^{23} CO molecules is _____. [1990]
 (A) 10 g molecules (B) 5 g molecules
 (C) 1 g molecules (D) 0.5 g molecules
- A 1 molar solution of H_2SO_4 is diluted from 1 litre to a volume of 10 litres, the normality of the solution will be _____. [1991]
 (A) 1 N (B) 0.1 N
 (C) 5 N (D) 0.5 N
- The amount of zinc required to produce 224 mL of H_2 at STP on treatment with dilute H_2SO_4 will be _____. [1996]
 (A) 65 g (B) 0.065 g
 (C) 0.65 g (D) 6.5 g

- In the reaction,
 $4NH_{3(g)} + 5O_{2(g)} \longrightarrow 4NO_{(g)} + 6H_2O_{(l)}$
 when 1 mole of ammonia and 1 mole of O_2 are made to react to completion _____. [1998]
 (A) all the oxygen will be consumed
 (B) 1.0 mole of NO will be produced
 (C) 1.0 mole of H_2O is produced
 (D) all the ammonia will be consumed
- Volume of CO_2 obtained by the complete decomposition of 9.85 g of $BaCO_3$ is _____. [2000]
 (A) 2.24 L (B) 2.24 L
 (C) 0.84 L (D) 0.84 L
- Molarity of liquid H_2O if density of solution is 1.17 g/cc is _____. [2001]
 (A) 36.5 (B) 18.25
 (C) 32.05 (D) 42.10
- What volume of oxygen gas (O_2) measured at $0^\circ C$ and 1 atm, is needed to burn completely 1 L of propane gas (C_3H_8) measured under the same conditions? [2008]
 (A) 5 L (B) 10 L
 (C) 7 L (D) 6 L
- How many moles of lead (II) chloride will be formed from a reaction between 6.5 g of PbO and 3.2 g HCl? [2008]
 (A) 0.011 (B) 0.029
 (C) 0.044 (D) 0.333
- 10 g of hydrogen and 64 g of oxygen were filled in a steel vessel and exploded. Amount of water produced in this reaction will be _____. [2009]
 (A) 3 mol (B) 4 mol
 (C) 1 mol (D) 2 mol
- 1.0 g of magnesium is burnt with 0.56 g O_2 in a closed vessel. Which reactant is left in excess and how much? (At wt. Mg = 24; O = 16) [2014]
 (A) Mg, 0.16 g (B) O_2 , 0.16 g
 (C) Mg, 0.44 g (D) O_2 , 0.28 g
- When 22.4 litres of $H_{2(g)}$ is mixed with 11.2 litres of $Cl_{2(g)}$, each at S.T.P, the moles of $HCl_{(g)}$ formed is equal to _____. [2014]
 (A) 1 mol of $HCl_{(g)}$
 (B) 2 mol of $HCl_{(g)}$
 (C) 0.5 mol of $HCl_{(g)}$
 (D) 1.5 mol of $HCl_{(g)}$
- 20.0 g of a magnesium carbonate sample decomposes on heating to give carbon dioxide and 8.0 g magnesium oxide. What will be the percentage purity of magnesium carbonate in the sample? (At. wt.: Mg = 24 u) [Re-Test 2015]
 (A) 60 (B) 84
 (C) 75 (D) 96



15. A mixture of 2.3 g formic acid and 4.5 g oxalic acid is treated with conc. H_2SO_4 . The evolved gaseous mixture is passed through KOH pellets. Weight (in g) of the remaining product at STP will be _____ [2018]
 (A) 1.4 (B) 3.0
 (C) 2.8 (D) 4.4

16. The number of moles of hydrogen molecules required to produce 20 moles of ammonia through Haber's process is _____. [2019]
 (A) 20 (B) 30
 (C) 40 (D) 10



Answers to MCQ's

- 1.1 : 1. (B)
 1.2 : 1. (D)
 1.3 : 1. (A) 2. (D)
 1.4 : 1. (D) 2. (A) 3. (C) 4. (D) 5. (B) 6. (A) 7. (D) 8. (A) 9. (C) 10. (A)
 11. (B) 12. (C) 13. (C) 14. (D) 15. (B) 16. (D) 17. (A) 18. (C) 19. (C)
 1.5 : 1. (C) 2. (A) 3. (B)
 1.6 : 1. (D) 2. (C) 3. (B) 4. (A) 5. (C) 6. (A) 7. (B) 8. (C) 9. (A) 10. (B)
 11. (B) 12. (A) 13. (A) 14. (B) 15. (C) 16. (B)



Hints to MCQ's

1.1 Units of measurement

1.

Quantity	Dimensions
Pressure	$[\text{M L}^{-1} \text{T}^{-2}]$
Force per unit volume	$[\text{M L}^{-2} \text{T}^{-2}]$
Energy per unit volume	$[\text{M L}^{-1} \text{T}^{-2}]$
Force	$[\text{M L T}^{-2}]$
Energy	$[\text{M L}^2 \text{T}^{-2}]$

1.2 Uncertainty in measurement

1. 161 has three significant figures as all are non-zero digits.
 0.161 has three significant figures as zero on the left of the first non-zero digit is not significant.
 0.0161 also has three significant figures as zeros on the left of the first non-zero digit are not significant.

1.3 Atomic and molecular masses

1. Average atomic mass

$$= \frac{\text{Sum of (Isotopic mass} \times \text{its abundance)}}{100}$$
 Average atomic mass = $\frac{(19 \times 10) + (81 \times 11)}{100}$
 $= 10.81 \approx 10.8$
2. Average atomic mass

$$= \frac{\text{Sum of (Isotopic mass} \times \text{its abundance)}}{100}$$
 Average isotopic mass of X

$$= \frac{(200 \times 90) + (199 \times 8) + (202 \times 2)}{100}$$

 $= 200 \text{ a.m.u.}$

1.4 Mole concept and molar mass

1. At STP,
 1 mol $\text{N}_2\text{O} = 22400 \text{ cc } \text{N}_2\text{O} = 6.02 \times 10^{23} \text{ } \text{N}_2\text{O}$ molecules
 $\therefore 1 \text{ cc of } \text{N}_2\text{O} = \frac{6.02 \times 10^{23}}{22400} \text{ molecules}$
 Each N_2O molecule contains 3 atoms,
 Hence,
 $\therefore 1 \text{ cc } \text{N}_2\text{O} = \frac{3 \times 6.02 \times 10^{23}}{22400} = \frac{1.8 \times 10^{22}}{22400}$
 Nitrogen contains 7 electrons while O contains 8 electrons. Hence, the number of electrons in one molecule of N_2O is 22.
 Hence,
 Number of electrons in 1 cc N_2O

$$= \frac{6.02 \times 10^{23}}{22400} \times 22 = \frac{1.32}{224} \times 10^{23} \text{ electrons}$$
2. Number of moles in 4.4 g of CO_2
 $= \frac{4.4}{44} = 0.1$
 Number of oxygen atoms in 1 mole of CO_2
 $= 2 \times N_A$
 \therefore Number of oxygen atoms in 0.1 mole of CO_2
 $= 0.1 \times 2 \times N_A$
 $= 0.2 \times 6.022 \times 10^{23}$
 $= 1.20 \times 10^{23}$
3. One litre of O_2 contains N molecules at 15°C and 150 mmHg pressure. If 1 L of one gas contains N molecules then 2 L of any gas under the same conditions will contain 2N molecules.



4. 1 L of air = $1000/0.21 = 210$ mL of O_2
 $\therefore 22400$ mL = 1 mole

$\therefore 210$ mL = $\frac{1}{22400} \times 210 = 0.0093$ mol

5. Weight of volatile gas = 0.24 g
 Volume of gas = 45 mL = 0.045 L

$$\text{Density} = \frac{\text{Mass}}{\text{Volume}}$$

$$\text{Mass of 45 mL of } H_2 = 0.089 \times 0.045 = 4.005 \times 10^{-3} \text{ g}$$

Vapour density

$$= \frac{\text{Mass of certain volume of vapour}}{\text{Mass of same volume of hydrogen}}$$

$$= \frac{0.24}{4.005 \times 10^{-3}} = 59.93$$

6. 100 g of haemoglobin contains 0.334 g of Fe

$\therefore 67200$ g of haemoglobin contains

$$= \frac{67200 \times 0.334}{100}$$

$$= 224.448 \text{ g of Fe.}$$

$$\text{Number of atoms of Fe} = \frac{224.448}{56}$$

$$= 4.008 \approx 4$$

7. Molecular mass of $NH_3 = 14 + (3 \times 1) = 17$

$$\text{Number of moles} = \frac{4.25}{17} = 0.25 \text{ mol}$$

Number of molecules of NH_3

$$= 0.25 \times 6.02 \times 10^{23} = 1.506 \times 10^{23} \text{ molecules}$$

One molecule of NH_3 contains 4 atoms.

$\therefore 1.506 \times 10^{23}$ molecules will contain

$$= 1.506 \times 10^{23} \times 4$$

$$= 6.024 \times 10^{23} \text{ atoms} \approx 6 \times 10^{23} \text{ atoms.}$$

8. Volume of cylindrical virus particle = $\pi r^2 l$

$$= 3.14 \times (7 \times 10^{-7})^2 \times 10^{-5} \times 10^3$$

$$= 1.54 \times 10^{-21} \text{ cc}$$

$$\text{Weight of the virus particle} = \frac{\text{Volume}}{\text{Specific volume}}$$

$$= \frac{1.54 \times 10^{-21}}{6.02 \times 10^{-2}}$$

\therefore Molecular weight of virus particle = weight of

$$1 \text{ } \mu\text{m}^3 \text{ particles} = \frac{1.54 \times 10^{-21}}{6.02 \times 10^{-2}} \times 6.02 \times 10^{23} \text{ g/mol}$$

$$= 15400 \text{ g/mol} = 15.4 \text{ kg/mol}$$

9. (A) 7 g $N_2 = \frac{7}{28} \times 6.022 \times 10^{23} = 1.51 \times 10^{23}$

(B) 2 g $H_2 = \frac{2}{2} \times 6.022 \times 10^{23} = 6.022 \times 10^{23}$

(C) 16 g $NO_2 = \frac{16}{46} \times 6.022 \times 10^{23} = 2.09 \times 10^{23}$

(D) 16 g $O_2 = \frac{16}{32} \times 6.022 \times 10^{23} = 2.26 \times 10^{23}$

10. (A) 15 L $H_2 = \frac{15}{22.4} \times 6.022 \times 10^{23} = 4.03 \times 10^{23}$

(B) 5 L $N_2 = \frac{5}{22.4} \times 6.022 \times 10^{23} = 1.34 \times 10^{23}$

(C) 0.5 g of $H_2 = \frac{0.5}{2} \times 6.022 \times 10^{23} = 1.51 \times 10^{23}$

(D) 10 g of $O_2 = \frac{10}{32} \times 6.022 \times 10^{23} = 1.88 \times 10^{23}$

11. Total number of atoms in a given amount of substance = $n \times N_A \times \text{Atomicity}$

$$= 0.1 \times 6.02 \times 10^{23} \times 3$$

$$= 1.806 \times 10^{23}$$

12. Option (A): 44 g $CO_2 = 1$ mole of CO_2

Option (B): 48 g $O_3 = 1$ mole of O_3

Option (C): 8 g $H_2 = 4$ moles of H_2

Option (D): 64 g $SO_2 = 1$ mole of SO_2

13. According to Avogadro's hypothesis, ratio of the volume of gases will be equal to the ratio of their no. of moles.

$$\frac{\text{weight of } H_2}{2} : \frac{\text{weight of } O_2}{32} : \frac{\text{weight of } CH_4}{16}$$

$$\frac{1}{2} : \frac{1}{2} : \frac{1}{16}$$

\therefore Ratio is $16 : 1 : 2$.

14. Number of moles of $H_2 = \frac{1}{2}$

$$\text{Number of moles of } O_2 = \frac{4}{32}$$

$$\text{Hence, molar ratio} = \frac{1}{2} : \frac{4}{32} = 4 : 1$$

15. 1 mole of water = 18 g of water

$$= 6.022 \times 10^{23} \text{ molecules of water}$$

$\therefore 18$ moles of water

$$= 18 \times 6.022 \times 10^{23} \text{ molecules of water}$$

$$= 1.08396 \times 10^{25} \text{ molecules of water}$$

16. When Avogadro number is $6.022 \times 10^{23} \text{ mol}^{-1}$, the mass of 1 mol of carbon = 12 g

\therefore Mass of 1 mol of carbon when Avogadro number is $6.022 \times 10^{20} \text{ mol}^{-1}$

$$= \frac{12 \times 6.022 \times 10^{20}}{6.022 \times 10^{23}} = 12 \times 10^{-3} \text{ g}$$

Thus, the mass of 1 mol of carbon is changed.

17. Volume of 1 mole of a gas at STP = 22.4 L

$$1 \text{ mol } CCl_4 \text{ vapour} = 12 + 4 \times 35.5$$

$$= 154 \text{ g}$$

Therefore, 22.4 L of a gas contains 154 g of CCl_4 .

$$\text{Density} = \frac{\text{Mass}}{\text{Volume}}$$

\therefore Density of CCl_4 vapour = $\frac{154}{22.4} \text{ g/L}$

$$= 6.875 \text{ g/L}$$



18. Option (A)

$$18 \text{ mL of water} = 18 \text{ g of water} = \frac{18}{18} \\ = 1 \text{ mol of water}$$

Option (B)

$$0.18 \text{ g water} = \frac{0.18}{18} = 0.01 \text{ mol of water}$$

Option (C)

$$0.00224 \text{ L of water vapours at 1 atm and} \\ 273 \text{ K (STP conditions)} = 2.24 \text{ mL of water} \\ = \frac{2.24}{22.4} \\ = 0.1 \text{ mol of water}$$

Option (D) has 10^{-3} mol of water.

Hence, 18 mL of water, i.e., option (A) has maximum number of moles of water and hence, it contains maximum number of water molecules.

19. Number of atoms

= Number of moles \times Avogadro's constant

$$= \frac{\text{Mass of a substance}}{\text{Molar mass of a substance}} \times N_A$$

$$\text{Number of atoms of Mg}_{(s)} = \frac{1}{24} \times N_A$$

$$\text{Number of atoms of O}_{2(g)} = \frac{2 \times 1}{32} \times N_A$$

$$\text{Number of atoms of Li}_{(s)} = \frac{1}{7} \times N_A$$

$$\text{Number of atoms of Ag}_{(s)} = \frac{1}{108} \times N_A$$

1.5 Percentage composition, empirical and molecular formulae

1. (A) % of nitrogen in $(\text{NH}_4)_2\text{SO}_4 = \frac{28}{132} \times 100$
21.21%

(B) % of nitrogen in $\text{C}_2\text{H}_7\text{N} = \frac{14}{40} \times 100$
35%

(C) % of nitrogen in $\text{CO}(\text{NH}_2)_2 = \frac{28}{60} \times 100$
46.66%

(D) % of nitrogen $\text{NH}_4\text{NO}_3 = \frac{28}{80} \times 100$
35%

0.5 mol of $\text{XY}_2 = 10 \text{ g}$

$$\therefore 1 \text{ mol of } \text{XY}_2 = 100 \text{ g}$$

i.e., Molecular weight of $\text{XY}_2 = 100$

0.05 mol of $\text{X}_3\text{Y}_2 = 9 \text{ g}$

$$\therefore 1 \text{ mol of } \text{X}_3\text{Y}_2 = 180 \text{ g}$$

i.e., Molecular weight of $\text{X}_3\text{Y}_2 = 180$

Let atomic weights of X and Y be x and y respectively.

$$\therefore x + 2y = 100 \quad \dots(i)$$

$$3x + 2y = 180 \quad \dots(ii)$$

Subtracting (i) from (ii),

$$2x = 180 - 100$$

$$\therefore x = 40$$

Substituting $x = 40$ in (i),

$$40 + 2y = 100$$

$$\therefore y = 30$$

3. Since, 0.5 g Se \equiv 100 gm peroxidase anhydrous enzyme

$$\therefore 78.4 \text{ g Se} = \frac{100 \times 78.4}{0.5} = 1.568 \times 10^4$$

Hence, minimum molecular mass of peroxidase anhydrous enzyme is $1.568 \times 10^4 \text{ g/mol}$.

1.6 Chemical reactions, stoichiometry and calculations based on stoichiometry

1. The reaction is



Hence, as per reaction stoichiometry, 1 mole H_2 or 2 g reacts with one mole of Z_2O_3 .

Now,

0.106 g of Z_2O_3 react with 0.006 g of H_2 .

$$\therefore 6 \text{ g } \text{H}_2 \text{ reacts with} = \frac{0.1596}{0.006} \times 6 = 159.6 \text{ g of}$$

Z_2O_3

Therefore, molecular mass of Z_2O_3 is 159.6 g/mol.

$$\therefore \text{Molecular mass of } \text{Z}_2\text{O}_3 = (2 \times \text{At. Wt. Z} + 3 \text{ At. Wt. O})$$

$$\text{Atomic mass of Z} = \frac{159.6 - (3 \times 16)}{2} = 55.8 \text{ g}$$

2. $\text{C}_2\text{H}_4 + 3\text{O}_2 \rightarrow 2\text{CO}_2 + 2\text{H}_2\text{O}$

28 g of ethylene require 96 g of O_2

$$\therefore 2.8 \times 10^3 \text{ g of ethylene require} = \frac{2.8 \times 10^3 \times 96}{28} \\ = 9.6 \times 10^3 \text{ g} \\ = 9.6 \text{ kg}$$

3. 1 mole of CO is equivalent to

$$6.02 \times 10^{23} \text{ molecules}$$

\therefore 10 mole CO will correspond to

$$6.02 \times 10^{24} \text{ molecules}$$

6.02×10^{24} CO molecules contain

6.02×10^{24} atoms of oxygen, which is equivalent to 10 g atoms of oxygen

10 g atoms of (O) oxygen = 5 g molecules of O_2 (\because Oxygen is a diatomic gas.)

4. $M_1V_1 = M_2V_2$

$$5 \times 1 = M_2 \times 10$$

$$M_2 = 0.5 \text{ M}$$

Normality = $n \times$ Molarity

$$= 2 \times 0.5$$

(\because H_2SO_4 is a diprotic acid)

$$= 1 \text{ N}$$



5. $\text{Zn} + \text{H}_2\text{SO}_4 \longrightarrow \text{ZnSO}_4 + \text{H}_2$
 1 Mole of zinc reacts to give 1 mole of hydrogen
 1 mole of hydrogen at STP is 22,400 mL.
 65 g zinc react to liberate 22400 mL of H_2
 \therefore Amount of zinc required to produce 224 mL of H_2 at STP = $\frac{224 \times 65}{22400} = 0.65 \text{ g}$

6. $4\text{NH}_3 + 5\text{O}_2 \longrightarrow 4\text{NO} + 6\text{H}_2\text{O}$
 From above reaction,
 4 Moles of NH_3 require 5 moles of O_2 .
 \therefore 1 Moles of $\text{NH}_3 = \frac{5}{4}$ moles of O_2
 $= 1.25 \text{ mol of O}_2$
 Therefore, 1 mol of NH_3 require 1.25 mol of O_2 . In given conditions, 1 mole of NH_3 and 1 mole of O_2 are made to react to completion. Hence, all the oxygen will be consumed.

7. $\text{BaCO}_3 \longrightarrow \text{BaO} + \text{CO}_2$
 197.34 g of BaCO_3 gives 22.4 L of CO_2
 \therefore 9.85 g of BaCO_3 will give $\frac{22.4 \times 9.85}{197.34}$
 $= 1.118 \text{ L} \approx 1.12 \text{ L}$

8. Density = 1.17 g/cc = 1170 g/L
 Hence, volume of the solution = 1 L
 Mass of the solute = 1170 g.
 Mol of solute = 32.05 mol
 Molarity = $\frac{\text{Moles of solute}}{\text{Volume of solution (L)}}$
 $= \frac{32.05}{1} = 32.05 \text{ M}$

9. $\text{C}_3\text{H}_8 + 5\text{O}_2 \longrightarrow 3\text{CO}_2 + 4\text{H}_2\text{O}$
 (1 mol). (5 mol)
 At STP, volume is proportional to mole.
 1 L of propane gas will require 5 L of O_2 .
 1 mol propane gas (C_3H_8) requires 5 mol oxygen gas (O_2). Hence, 1 L propane gas (C_3H_8) requires 5 L oxygen gas (O_2).

10. $\text{PbO} + 2\text{HCl} \longrightarrow \text{PbCl}_2 + \text{H}_2\text{O}$
 Molecular weight of $\text{PbO} = 207.2 + 16$
 $= 223.2$
 Moles of $\text{PbO} = \frac{6.5}{223.2} = 0.029 \text{ mol}$
 Moles of $\text{HCl} = \frac{3.2}{36.5} = 0.088 \text{ mol}$
 0.029 mol of PbO required 0.058 mol of HCl .
 Hence, HCl is in excess, PbO is limiting reagent.
 From stoichiometry, mol of $\text{PbO} = \text{mol of PbCl}_2$
 0.029 mol of $\text{PbO} = 0.029 \text{ mol of PbCl}_2$

11. $2\text{H}_{2(\text{g})} + \text{O}_{2(\text{g})} \longrightarrow 2\text{H}_2\text{O}_{(\text{g})}$
 Ratio of moles of reactants, $\text{H}_2 : \text{O}_2 = 2 : 1$
 Actual amount of reactants: 10 g H_2 and 64 g O_2
 Actual moles of reactants: 5 mol H_2 and 2 mol O_2
 Ratio of actual moles of reactants,
 $\text{H}_2 : \text{O}_2 = 5 : 2 = 2.5 : 1$

\therefore The limiting reactant is O_2 .
 Now, 1 mole of oxygen gives 2 moles of water. Hence, 2 moles of oxygen will give 4 moles of water.

12. $2\text{Mg} + \text{O}_2 \longrightarrow 2\text{MgO}$
 $(2 \times 24) \quad (32)$
 48 g of Mg require 32 g of O_2
 \therefore 0.56 g of O_2 require $\frac{0.56 \times 48}{32}$
 $= 0.84 \text{ g of Mg}$
 \therefore $\text{Mg left} = 1 - 0.84 = 0.16 \text{ g}$

13. 1 mol gas \equiv 22.4 L at S.T.P.
 Moles of $\text{Cl}_2 = \frac{11.2}{22.4} = 0.5 \text{ mol}$
 The reaction is
 $\text{H}_{2(\text{g})} + \text{Cl}_{2(\text{g})} \longrightarrow 2\text{HCl}_{(\text{g})}$
 From the reaction, 1 mol of H_2 requires 1 mol of Cl_2 to form 2 mol of HCl . Since, available Cl_2 is 0.5 mol, it is limiting reactant.
 \therefore Hence, 1 mol $\text{Cl}_2 = 2 \text{ mol HCl}$
 0.5 mol $\text{Cl}_2 = 1 \text{ mol HCl}$

14. $\text{MgCO}_{3(\text{s})} \longrightarrow \text{MgO}_{(\text{s})} + \text{CO}_{2(\text{g})}$
 Molar mass of $\text{MgCO}_3 = 84 \text{ g mol}^{-1}$
 \therefore Number of moles of $\text{MgCO}_3 = \frac{20}{84} = 0.238 \text{ mol}$
 \therefore 1 mole MgCO_3 gives 1 mole MgO
 \therefore 0.238 mole MgCO_3 will give 0.238 mole MgO .
 Molar mass of $\text{MgO} = 40 \text{ g mol}^{-1}$
 \therefore 0.238 mole $\text{MgO} = 40 \times 0.238$
 $= 9.52 \text{ g MgO}$
 \therefore Theoretical yield of $\text{MgO} = 9.52 \text{ g}$
 Practical yield of MgO is 8.0 g
 \therefore Percentage purity = $\frac{8}{9.52} \times 100 = 84 \%$


15. $\text{HCOOH} \xrightarrow{\text{Conc. H}_2\text{SO}_4} \text{CO} + \text{H}_2\text{O}$
 0.5 mol 0.5 mol
 $(\text{COOH})_2 \xrightarrow{\text{Conc. H}_2\text{SO}_4} \text{CO} + \text{CO}_2 + \text{H}_2\text{O}$
 0.5 mol 0.5 mol
 Gaseous mixture formed is CO and CO_2 . When it is passed through KOH , which CO_2 is absorbed. So, the remaining gas is CO .
 Weight of remaining gaseous product at STP is $0.5 \times 0.5 \times 28 = 2.8 \text{ g}$

16. $\text{N}_{2(\text{g})} + 3\text{H}_{2(\text{g})} \longrightarrow 2\text{NH}_{3(\text{g})}$
 3 mol $\text{H}_2 = 2 \text{ mol NH}_3$
 \therefore 30 mol $\text{H}_2 = 20 \text{ mol NH}_3$

NEET



PSP PREVIOUS SOLVED PAPERS



AVAILABLE SUBJECTS:

NEET Physics

NEET Chemistry

NEET Biology

BUY NOW

Salient Features

- Important questions with solutions from 1988 to 2020 (33 years of AIPMT and 8 years of NEET)
- Graphical Analysis of subtopics at the beginning renders detailed insight about subtopic marks distribution
- Includes 1571 AIPMT/ NEET MCQs to foster better preparation
- Includes Solved questions from NEET 2020 to polish strengthen concept base



Target Publications® Pvt. Ltd.
Transforming lives through learning.

88799 39712 / 13 / 14 / 15

mail@targetpublications.org

www.targetpublications.org